



Plant Poisonings in Australia; A Review of Calls to the Queensland Poisons Information Centre



Islam T², Knoeckel R^{1,2}, Wylie C¹ and Isoardi KZ.^{1,2}
1 Queensland Poisons Information Centre, Queensland Children's Hospital, Brisbane, Australia.
2 Faculty of Medicine, University of Queensland, Brisbane, Australia.



Introduction

- Plant exposures are a common reason for people to seek advice from Poisons Information Centres.¹
- The literature describing plant poisonings in Australia is very limited and focusses only on exposures resulting in significant toxicity.
- We aimed to characterise all plant exposures referred to an Australian state-based Poisons Information Centre, with a particular focus on symptomatology and referral advice

Methods

- This study is a retrospective review of calls to the Queensland Poisons Information Centre (QPIC) regarding plant exposures between January 2018 – December 2021
- Data was extracted from the QPIC for patient demographics (age, sex), caller details (site, relation to patient), exposure characteristics (plant, site of exposure, route of exposure, any co-ingestions, symptoms) and referral advice (non-medical or medical).
- Chart review was performed for severe cases.
- Poison Severity Scores (PSS) for acute poisoning were calculated for the severity of poisoning in each case.²
- Descriptive statistics were used, and performed in GraphPad Prism 7.0d.

Results

- Overall, there were 3155 calls (including 389 recalls) recorded.
- The number of plant-related calls increased each year, from 810 in 2019 to 1003 in 2021.
- Most calls were made concerning exposures in children aged 1-4 years (46.8%). These were most frequently made by a family member or carer (73.5%). Most calls were from home (81.6%).
- Plant exposures were mostly unintentional (98.4%), and the oral route of exposure was most common (81.9%).
- The plant involved was known in 2366 calls (85.5%). The commonest plant groups are graphed below:



- By species, the most frequent plants exposed were *Euphorbia spp.* (9.6%), Elephant's Ear (6.5%), and *Duranta Erecta* (5.6%)
- Patients were mostly asymptomatic (59.4%) or experienced mild poisoning (37.4%). Only 18 (0.6%) experienced moderate or severe poisonings.
- Severe poisonings was mostly observed in male adults using plants for their sedative or anticholinergic properties.
- Medical review was advised in 409 (12.9%) of cases. These were mostly in adults following ocular exposures to *Euphorbiaceae*.

Discussion & Conclusions

- Plant exposures were mostly accidental paediatric exposures and relatively benign. In the majority of cases, patients were either asymptomatic or experienced mild toxicity that required no medical referral.
- Medical review was advised more commonly following exposures to *Euphorbiaceae*, gastrointestinal irritants, and oxalates. Severe toxicity was rare, and most often occurred in young adults using plants for recreational purposes.
- The findings of our study are consistent with the wider literature reporting on plant poisonings globally.³⁻⁴

References

1. Queensland Poisons Information Centre. Annual Report 2019. Brisbane: Queensland Children's Hospital; 2019.
2. Casey P, Dexter E, Michell J, Vale J. The Prospective Value of the IPCS/EC/EAPCCT Poisoning Severity Score in Cases of Poisoning. *Journal of Toxicology: Clinical Toxicology*. 1998;36(3):215-217
3. Krenzelok E, Mrvos R. Friends and foes in the plant world: A profile of plant ingestions and fatalities. *Clinical Toxicology*. 2011;49(3):142-149
4. Fuchs J, Rauber-Lüthy C, Kupferschmidt H, Kupper J, Kullak-Ublick G, Ceschi A. Acute plant poisoning: Analysis of clinical features and circumstances of exposure. *Clinical Toxicology*. 2011;49(7):671-680.